

Isgur-Wise Function for $\Lambda_b \rightarrow \Lambda_c$ in B-S Approach

X.-H. Guo^{1,2} and T. Muta¹

1. Department of Physics, Hiroshima University, Japan

2. Institute of High Energy Physics, Academia Sinica, Beijing, China

Abstract

In the heavy quark limit, the heavy baryon Λ_Q ($Q=b$ or c) can be regarded as composed of a heavy quark and a scalar light diquark which has good spin and flavor quantum numbers. Based on this picture we establish the Bethe-Salpeter (B-S) equation for Λ_Q in the leading order of $1/m_Q$ expansion. With the kernel containing both the scalar confinement and one-gluon-exchange terms we solve the B-S equation numerically. The Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ is obtained numerically from our model. Comparison with other model calculations are also presented. It seems that the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ drops faster than that for $B \rightarrow D$. The differential and total decay widths for $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$ are given in the limit $m_{b,c} \rightarrow \infty$.

PACS Numbers: 11.10.St, 12.39.Hg, 14.40.Nd, 14.40.Lb

I. Introduction

The past few years have seen much progress in heavy flavor physics due to the discovery of the new flavor and spin symmetries $SU(2)_f \times SU(2)_s$ in the heavy quark limit and the establishment of the heavy quark effective theory (HQET) [1]. In the framework of the HQET the processes involving heavy quarks can be simplified since the HQET may reduce the number of form factors. For instance in the leading order of $1/m_Q$ expansion only one form factor (the Isgur-Wise function) remains for $\Lambda_b \rightarrow \Lambda_c$ transition. However, to get the complete physics one still has to face the basic problem, the nonperturbative effects of QCD. As the formally exact equation to describe the relativistic bound state, the B-S equation [2][3] can be simplified to a great extent in the heavy quark limit and has been applied to give many theoretical results concerning heavy mesons [4]. At present, there are not many experimental results for heavy baryons. But in the future we may expect more and more data coming out in LEP and the forthcoming B-factory. Actually recently OPAL has measured some physical quantities for Λ_b such as the lifetime of Λ_b and the product branching ratio for $\Lambda_b \rightarrow \Lambda l^- \bar{\nu} X$ [5]. Hence the study of heavy baryons are of particular interest in the near future. It is the motivation of the present paper to establish the B-S equation for the heavy baryon in the heavy quark limit and then solve it numerically by assuming some reasonable form of its kernel, and consequently give some phenomenological predictions.

When the quark mass is very heavy comparing with the QCD scale Λ_{QCD} , the light degrees of freedom in a heavy baryon Λ_Q ($Q=b$ or c) becomes blind to the flavor and spin quantum numbers of the heavy quark because of the $SU(2)_f \times SU(2)_s$ symmetries. Therefore, the angular momentum and flavor quantum numbers of the light degrees of freedom (the light diquark) become good quantum numbers which can be used to classify heavy baryons. For example, Λ_Q and $\Sigma_Q^{(*)}$ correspond to the zero and one angular momentum states of the light diquark respectively. Hence

it is natural to regard the heavy baryon to be composed of a heavy quark and a light diquark. In the present paper we concentrate on Λ_Q which has a scalar light diquark $S_{[ud]}$ with $[ud]$ flavor quantum number and zero spin and isospin. Other heavy baryons like $\Sigma_Q^{(*)}$ can be dealt with in the similar way.

Based on the above picture of the composition of the heavy baryon the three body system is simplified to two body system. We will establish the B-S equation for the heavy baryon in this picture. Then we will solve this equation numerically by assuming that the kernel contains the scalar confinement and one-gluon-exchange terms. Then we will apply the B-S equation to calculate the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ and the decay width for the semileptonic decay of Λ_b to Λ_c in the order $m_{b,c} \rightarrow \infty$.

The remainder of this paper is organized as follows. In Sect. II we establish the B-S equation for the heavy quark and light scalar diquark system and discuss the form of its kernel. Consequently this equation can be solved numerically. Then in Sect. III we apply the solution of the B-S equation to calculate the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ and also the differential and total decay widths for $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$. Comparison with other model calculations for the Isgur-Wise function will also be presented. Finally, Sect. VI is served for summary and discussions.

II. The B-S equation for Λ_Q

As discussed in Introduction Λ_Q is regarded as the bound state of a heavy quark and a light scalar diquark. Based on this picture we can define the B-S wave function of Λ_Q by

$$\chi(x_1, x_2, P) = \langle 0 | T \psi(x_1) \varphi(x_2) | \Lambda_Q(P) \rangle, \quad (1)$$

where $\psi(x_1)$ and $\varphi(x_2)$ are the wave functions of the heavy quark and the light scalar diquark respectively, $P = m_{\Lambda_Q} v$ is the momentum of Λ_Q and v is its velocity. Let m_Q and m_D be the masses of the heavy quark and the light diquark in the baryon,

$\lambda_1 = \frac{m_Q}{m_Q+m_D}$, $\lambda_2 = \frac{m_D}{m_Q+m_D}$, p be the relative momentum of the two constituents.

The B-S wave function in the momentum space is defined as

$$\chi(x_1, x_2, P) = e^{iPX} \int \frac{d^4p}{(2\pi)^4} e^{ipx} \chi_P(p), \quad (2)$$

where $X = \lambda_1 x_1 + \lambda_2 x_2$ is the coordinate of the center of mass and $x = x_1 - x_2$.

The momentum of the heavy quark is $p_1 = \lambda_1 P + p$ and that of the diquark is $p_2 = -\lambda_2 P + p$.

The derivation of the B-S equation for the two fermion systems can be found in textbooks [3]. In the same way we can prove that for the fermion and scalar object system the form of the B-S equation is still valid. $\chi_P(p)$ satisfies the following B-S equation

$$\chi_P(p) = S_F(\lambda_1 P + p) \int \frac{d^4q}{(2\pi)^4} G(P, p, q) \chi_P(q) S_D(-\lambda_2 P + p), \quad (3)$$

where $G(P, p, q)$ is the kernel which is defined as the sum of the two particle irreducible diagrams. In the following we will use the variables $p_l = v \cdot p - \lambda_2 m_{\Lambda_Q}$, $p_t = p - (v \cdot p)v$. Then in the leading order of $1/m_Q$ expansion we have

$$\begin{aligned} S_F(\lambda_1 P + p) &= \frac{i(1 + \not{v})}{2(p_l + E_0 + m_D + i\epsilon)}, \\ S_D(-\lambda_2 P + p) &= \frac{i}{p_t^2 - W_p^2 + E_0 + i\epsilon}, \end{aligned} \quad (4)$$

where $W_p = \sqrt{p_t^2 + m_D^2}$ and E_0 is the binding energy. The corrections to eq. (4) are from $O(1/m_Q)$.

In general, $\chi_P(p)$ can be expanded as the following form,

$$\chi_P(p) = (A + B\not{v} + C\not{p} + D\not{p}\not{p})u_{\Lambda_Q}(v, s), \quad (5)$$

where $u_{\Lambda_Q}(v, s)$ is the spinor of Λ_Q with helicity s and A, B, C, D are Lorentze scalar functions.

From eqs. (3) and (4) it can be seen that

$$\not{p}\chi_P(p) = \chi_P(p). \quad (6)$$

Combining eqs. (5) and (6) we immediately see that

$$\chi_P(p) = \phi_P(p)u_{\Lambda_Q}(v, s), \quad (7)$$

where $\phi_P(p)$ is a scalar field. This form is consistent with our picture about the heavy baryon.

We assume the kernel has the form

$$-iG = I \otimes IV_1 + v_\mu \otimes (p_2 + p'_2)^\mu V_2, \quad (8)$$

where the first term arises from scalar confinement and the second one is from one gluon exchange diagram. The vertex of gluon with two scalar diquarks is proportional to $(p_2 + p'_2)_\mu$ (p_2 and p'_2 are the momenta of the two scalar diquarks on the vertex) and a form factor $F((p_2 + p'_2)^2)$ (see Fig. 1). Substituting eqs. (4), (7) and (8) into (3) we have

$$\phi_P(p) = -\frac{1}{(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^4q}{(2\pi)^4} G(P, p, q) \phi_P(q). \quad (9)$$

Now consider the vertex of two heavy quarks with gluon. The momenta of the two heavy quarks are $p_1 = \lambda_1 m_{\Lambda_Q} v + p$ and $p'_1 = \lambda_1 m_{\Lambda_Q} v + q$ respectively. p and q are of the order Λ_{QCD} . In the heavy quark limit the heavy quark is almost on-shell, therefore, $p_l = q_l$ on this vertex. Hence we can make the covariant instantaneous approximation [4] in the kernel. Then eq. (9) becomes

$$\phi_P(p) = -\frac{i}{(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^4q}{(2\pi)^4} (\tilde{V}_1 + 2p_l \tilde{V}_2) \phi_P(q), \quad (10)$$

where $\tilde{V}$ is defined as $\tilde{V} \equiv V|_{p_l=q_l}$.

In general $\phi_P(p)$ can be the function of p_l and p_t . Defining $\tilde{\phi}_P(p_t) = \int \frac{dp_l}{2\pi} \phi_P(p)$ one gets immediately the B-S equation for $\tilde{\phi}_P(p_t)$

$$\tilde{\phi}_P(p_t) = -\frac{1}{2(E_0 - W_p + m_D)W_p} \int \frac{d^3 q_t}{(2\pi)^3} (\tilde{V}_1 - 2W_p \tilde{V}_2) \tilde{\phi}_P(q_t). \quad (11)$$

If one knows the form for the kernel $\tilde{V}_1$ and $\tilde{V}_2$, then $\tilde{\phi}_P(p_t)$ can be obtained. Consequently $\phi_P(p)$ can be solved since after integrating q_l in eq. (10) we get the relation between $\phi_P(p)$ and $\tilde{\phi}_P(q_t)$

$$\phi_P(p) = \frac{i}{(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^3 q_t}{(2\pi)^3} (\tilde{V}_1 + 2p_l \tilde{V}_2) \tilde{\phi}_P(q_t). \quad (12)$$

The kernel $\tilde{V}_1$ and $\tilde{V}_2$ for the B-S equation in the meson case was given in [4] as

$$\tilde{V}_1|_{\text{meson}} = \frac{8\pi\kappa'}{[(p_t - q_t)^2 + \mu^2]^2} - (2\pi)^3 \delta^3(p_t - q_t) \int \frac{d^3 k}{(2\pi)^3} \frac{8\pi\kappa'}{(k^2 + \mu^2)^2}, \quad (13)$$

$$\tilde{V}_2|_{\text{meson}} = -\frac{16\pi}{3} \frac{\alpha_{seff}}{(p_t - q_t)^2 + \mu^2}, \quad (14)$$

where κ' and α_{seff} are coupling parameters related to scalar confinement and one-gluon-exchange diagram respectively. The second term in eq. (13) is the counter term which removes the infra-red divergence in the integral equation. From potential model κ' is from 0.18 to 0.2. The parameter μ is introduced to avoid the infra-red divergence in numerical calculations. The limit $\mu \rightarrow 0$ is taken in the end.

In the baryon case, since confinement is still due to scalar interaction the form of $\tilde{V}_1$ need not be changed. Only the parameter κ' has to be replaced by κ which describes the confinement interaction between the quark and diquark. However, since the diquark is not a point-like object, there should be a form factor $F(Q^2)$ ($Q = p_2 - p'_2$) to describe the vertex of the gluon and two scalar diquarks. In general, this vertex is (Fig. 1)

$$ig_s \frac{\lambda^a}{2} (p_2 + p'_2)^\mu F(Q^2)$$

with λ 's being the Gell-Mann color matrices and g_s the strong coupling constant. The form of $F(Q^2)$ was assumed as [6]

$$F(Q^2) = \frac{\alpha_{seff} Q_0^2}{Q^2 + Q_0^2}, \quad (15)$$

where Q_0^2 is a parameter which freezes $F(Q^2)$ when Q^2 is very small. In the high energy region the form factor is proportional to $1/Q^2$ which is consistent with perturbative QCD calculations [7]. By analyzing the eletromagnetic form factor for the proton it was found that $Q_0^2 = 3.2\text{GeV}^2$ can lead to consistent results with the experimental data [6].

Based on the above analysis the kernel for the B-S equation in the baryon case is of the following form

$$\begin{aligned} \tilde{V}_1 &= \frac{8\pi\kappa}{[(p_t - q_t)^2 + \mu^2]^2} - (2\pi)^3 \delta^3(p_t - q_t) \int \frac{d^3k}{(2\pi)^3} \frac{8\pi\kappa}{(k^2 + \mu^2)^2}, \\ \tilde{V}_2 &= -\frac{16\pi}{3} \frac{\alpha_{seff}^2 Q_0^2}{[(p_t - q_t)^2 + \mu^2][(p_t - q_t)^2 + Q_0^2]}. \end{aligned} \quad (16)$$

There are two parameters κ and α_{seff} in the kernel. However, there should some relation between them since when we solve the B-S equation (11) numerically the binding energy should satisfy the following relation

$$m_{\Lambda_Q} = m_Q + m_D + E_0, \quad (17)$$

where we have omitted corrections from $O(1/m_Q)$ since we are working in the heavy quark limit. From the B-S equation solutions in meson case it has been found that the values $m_b = 5.02\text{GeV}$ and $m_c = 1.58\text{GeV}$ give predictions which are in good agreement with experiments [4]. Hence in the baryon case we expect

$$m_D + E_0 = 0.62\text{GeV}. \quad (18)$$

On the other hand the dimesion of κ is three and that of κ' is two. This extra dimesion in κ should be caused by nonperturbative diagrams which include the

frozen form factor $F(Q^2)$ at low momentum region. Since Λ_{QCD} is the only parameter which is related to confinement we expect that

$$\kappa \sim \Lambda_{QCD} \kappa'. \quad (19)$$

Therefore in our numerical calculations we let κ vary in the region between 0.02GeV^3 to 0.1GeV^3 . The diquark mass m_D is chosen to vary from 650 MeV to 800 MeV. In order to satisfy the relation (18) we obtain the parameters for different values of m_D . The results for $m_D = 700\text{MeV}$ are shown in Table 1.

Table 1. Values of κ and α_{seff} for $m_D = 700\text{MeV}$

$\kappa(\text{GeV}^3)$	0.02	0.04	0.06	0.08	0.1
α_{seff}	0.67	0.70	0.72	0.74	0.75

When $m_D = 650\text{MeV}$ we find that $\alpha_{seff} = 0.60$ for $\kappa = 0.02\text{GeV}^3$ and $\alpha_{seff} = 0.71$ for $\kappa = 0.1\text{GeV}^3$. When $m_D = 800\text{MeV}$, $\alpha_{seff} = 0.79$ for $\kappa = 0.02\text{GeV}^3$ and $\alpha_{seff} = 0.84$ for $\kappa = 0.1\text{GeV}^3$. Having these two parameters we obtain the numerical solution for the B-S wave function directly.

III. The Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$

In this section we will apply the B-S equation for heavy baryons to obtain the numerical result for the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$. The transition diagram is plotted in Fig.2. In the limit $m_{b,c} \rightarrow \infty$ there is only one form factor, the Isgur-Wise function to describe the weak transition from Λ_b to Λ_c ,

$$\langle \Lambda_c(v') | \bar{c} \gamma_\mu b | \Lambda_b(v) \rangle = \xi(\omega) \bar{u}_{\Lambda_c}(v') \gamma_\mu u_{\Lambda_b}(v), \quad (20)$$

where $\omega = v \cdot v'$ is the velocity transfer and u_{Λ_b} and u_{Λ_c} are the Dirac spinors of Λ_b and Λ_c respectively.

On the other hand the transition matrix element of $\Lambda_b \rightarrow \Lambda_c$ is related to the B-S wave functions of Λ_b and Λ_c by the following equation

$$\langle \Lambda_c(v') | \bar{c} \gamma_\mu b | \Lambda_b(v) \rangle = \int \frac{d^4 p}{(2\pi)^4} \bar{\chi}_{P'}(p') \chi_P(p) S_D^{-1}(p_2), \quad (21)$$

where P (P') is the momentum of Λ_b (Λ_c). $\bar{\chi}_{P'}(p')$ is the wave function of the final state $\Lambda_c(v')$ which satisfies the constraint

$$\bar{\chi}_{P'}(p) \not{p}' = \bar{\chi}_{P'}(p). \quad (22)$$

The scalar part of the final state B-S wave function obeys the same B-S equation as (9). Then from eqs. (20) and (21) one gets immediately

$$\xi(\omega) = \int \frac{d^4 p}{(2\pi)^4} \phi_{P'}(p') \phi_P(p) S_D^{-1}(p_2). \quad (23)$$

Since in the weak transition the diquark acts as a spectator its momentum in the initial and final baryons should be the same, $p_2 = p'_2$. Then we can show that

$$p' = p + m_D(v' - v), \quad (24)$$

where again we omitted the $O(1/m_Q)$ corrections. From eq. (24) we can get the relations between p'_l , p'_t and p_l , p_t straightforwardly

$$\begin{aligned} p'_l &= p_l \omega - p_t \sqrt{\omega^2 - 1} \cos \theta, \\ p'^2_t &= p_t^2 + p_t^2 (\omega^2 - 1) \cos^2 \theta + p_l^2 (\omega^2 - 1) - 2 p_l p_t \omega \sqrt{\omega^2 - 1} \cos \theta, \end{aligned} \quad (25)$$

where θ is the angle between p_t and v'_t .

Substituting the relation between $\phi_P(p)$ and $\tilde{\phi}_P(p_t)$ (12) into eq. (23) and after integrating the p_l component by selecting the proper contour we have

$$\begin{aligned} \xi(\omega) &= \int \frac{d^3 p_t}{(2\pi)^3} \frac{1}{2W_p(E_0 + m_D - W_p)(E_0 + m_D - \omega W_p - p_t \sqrt{\omega^2 - 1} \cos \theta)} \\ &\cdot \int \frac{d^3 r_t}{(2\pi)^3} [\tilde{V}_1(p'_t - r_t) - 2(\omega W_p + p_t \sqrt{\omega^2 - 1} \cos \theta) \tilde{V}_2(p'_t - r_t)] |_{k_l = -W_p} \tilde{\phi}_{P'}(r_t) \\ &\cdot \int \frac{d^3 l_t}{(2\pi)^3} [\tilde{V}_1(p_t - l_t) - 2\omega W_p \tilde{V}_2(p_t - l_t)] \tilde{\phi}_P(l_t), \end{aligned} \quad (26)$$

where the relation (25) has been substituted into the above equation.

In eq. (26) all the three-dimensional integral can be simplified to one-dimensional integral. Furthermore, the normalization constant of the B-S wave function should be chosen such that the Isgur-Wise function is equal to one at zero momentum transfer. In Fig. 3 we plot numerical result for the Isgur-Wise function with the parameter $m_D = 700$ MeV. For other values of m_D in the range of 650MeV to 800MeV the shape of the Isgur-Wise function does not change a lot.

The slope of the Isgur-Wise function at $\omega = 1$ which is of particular interests is defined as

$$\frac{d\xi(\omega)}{d\omega} \big|_{\omega=1} = -\rho^2. \quad (27)$$

In Table 2 we list the values of ρ^2 for different m_D and κ . The values of ρ^2 outside (inside) the braket correspond to $\kappa = 0.02\text{GeV}^3$ (0.1GeV^3) respectively.

Table 2. Values of ρ^2

$m_D(\text{GeV})$	0.65	0.7	0.8
ρ^2	1.4 (2.4)	1.4 (2.4)	1.6 (2.4)

It can be seen from Table 2 that the slope is insensitive to the values of m_D .

The Isgur-Wise function has also been calculated in other models. In Ref. [8] Guo and Kroll use the Drell-Yan type overlap integrals for the model hadronic wave functions of Λ_b and Λ_c to obtain the following form of the Isgur-Wise function

$$\xi(\omega) = \left(\frac{2}{\omega + 1} \right) \exp \left(-2\varepsilon^2 b^2 \frac{\omega - 1}{\omega + 1} \right) \frac{K_6(2\varepsilon b / \sqrt{\omega + 1})}{K_6(\sqrt{2}\varepsilon b)}, \quad (28)$$

where ε is the light scalar diquark mass and b is related to the average transverse momentum of the constituents in the heavy baryon. K_l is defined as

$$K_l(x) = \int_{-x}^{\infty} dz e^{-z^2} (z + x)^l.$$

The Isgur-Wise function obtained by Jenkins, Manohar and Wise [9] from the soliton model has the following form

$$\xi(\omega) = 0.99 \exp[-1.3(\omega - 1)]. \quad (29)$$

The MIT bag model calculation by Sadzikowski and Zalewski [10]) gives the following result

$$\xi(\omega) = \left(\frac{2}{\omega + 1} \right)^{3.5 + 1.2/\omega}. \quad (30)$$

The values of ρ^2 from the above models are listed in the following

$$\rho^2 = \begin{cases} 2.9(3.7)[8] & \text{for } b = 1.18(1.77) \text{ GeV}^{-1}, \\ 1.3[9], \\ 2.4[10]. \end{cases} \quad (31)$$

We can see that different models may give results for the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ with some difference. $\rho^2 \geq 1.3$ in all these model calculations. The ρ parameter of the Isgur-Wise function for $B \rightarrow D$ has also been calculated. In the B-S equation approach [4] ρ^2 varies from 1.0 to 1.1. Other model calculations for ρ in $B \rightarrow D$ transition gives the following values: $\rho^2 = 1.19 \pm 0.25[11]$; $\rho = 1.13 \pm 0.11[12]$; and $\rho = 1.20 \pm 0.17[13]$. Because of the large uncertainties we can not draw a definite conclusion by comparing the slopes of the Isgur-Wise function at the zero recoil point in the meson and baryon cases. However, it seems that the Isgur-Wise function drops faster in the baryon case than in the meson case. At least in the B-S approach the ρ parameter is bigger in the baryon case than in the meson case.

The decay width for the semileptonic transition $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$ can also be predicted by using the Isgur-Wise function obtained. The differential decay width has the following form [8] [14]

$$\frac{d\Gamma}{d\omega} = \frac{2}{3} m_{\Lambda_c}^4 m_{\Lambda_b} A \xi^2(\omega) \sqrt{\omega^2 - 1} [3\omega(\eta + \eta^{-1}) - 2 - 4\omega^2], \quad (32)$$

where $\eta = m_{\Lambda_b}/m_{\Lambda_c}$ and $A = \frac{G_F^2}{(2\pi)^3} |V_{cb}|^2 B(\Lambda_c \rightarrow ab)$. $|V_{cb}|$ is the Kobayashi-Maskawa matrix element. $B(\Lambda_c \rightarrow ab)$ is the branching ratio for the decay $\Lambda_c \rightarrow a(\frac{1}{2}^+) + b(0^-)$

through which Λ_c is detected since the structure for such decay is already well known. The plot for $A^{-1}\frac{d\Gamma}{d\omega}$ is shown in Fig. 4 for $m_D = 700\text{MeV}$. Again for other values of m_D the results change only a little.

After integrating ω in eq. (32) we have the total decay width for $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$. For $m_D = 700\text{MeV}$

$$\Gamma(\Lambda_b \rightarrow \Lambda_c l \bar{\nu}) = \begin{cases} 4.2B(\Lambda_c \rightarrow ab) \times 10^{10}s^{-1} & \text{when } \kappa = 0.02\text{GeV}^3, \\ 5.7B(\Lambda_c \rightarrow ab) \times 10^{10}s^{-1} & \text{when } \kappa = 0.1\text{GeV}^3. \end{cases} \quad (33)$$

When $m_D = 650\text{MeV}$, $\Gamma(\Lambda_b \rightarrow \Lambda_c l \bar{\nu})$ is $4.3(5.9)B(\Lambda_c \rightarrow ab) \times 10^{10}s^{-1}$ for $\kappa = 0.02(0.1)$ and when $m_D = 800\text{MeV}$, $\Gamma(\Lambda_b \rightarrow \Lambda_c l \bar{\nu})$ is $4.1(5.0)B(\Lambda_c \rightarrow ab) \times 10^{10}s^{-1}$ for $\kappa = 0.02(0.1)$.

IV. Summary and discussions

In this paper we established the B-S equation for the heavy baryon which is considered as composed of a heavy quark and a light scalar diquark. By assuming the kernel containing scalar confinement and one-gluon-exchange terms we solve the B-S equation numerically. Furthermore, we applied the obtained results to calculate the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$. It is found that in the B-S approach, the Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c$ decreases faster than for $B \rightarrow D$. Comparison with other model calculations are also presented. The slope parameter ρ^2 at the zero recoil point for $\Lambda_b \rightarrow \Lambda_c$ is larger than 1.3 in all the present model calculations. In the limit $m_{b,c} \rightarrow \infty$ we also calculated the differential and total decay widths for $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$.

Although the B-S equation is formally the exact equation to describe the bound state, there is much difficulty in applying it to the real physical state. The most difficult point is the that we can not solve out the form of the kernel. Hence we have to use some phenomenological kernel which is reasonable because of the success of the potential model. This leads to some uncertainties. In our approach, the parameters κ and α_{seff} in the kernel are not exactly determined. Therefore, we let

them to vary in some reasonable range. The experimental data concerning Λ_Q in the future can help to fix the parameters in our model.

We have worked in the heavy quark limit. The physical predictions such as $\Gamma(\Lambda_b \rightarrow \Lambda_c l \bar{\nu})$ will be subjective to $1/m_Q$ corrections. The study on the $1/m_Q$ corrections will appear in our later work.

Acknowledgment:

One of us (X.-H. G) would like to thank the Japan Society for the Promotion of Science (JSPS) for the support. He is also indebted to Dr. H.-Y. Jin for helpful discussions.

References

- [1] N. Isgur and M.B. Wise, Phys. Lett. **B232** (1989) 113, **B237** (1990) 527; H. Georgi, Phys. Lett. **B264** (1991) 447; see also M. Neubert, Phys. Rep. **245** (1994) 259 for the review.
- [2] E.E. Salpeter and H.A. Bethe, Phys. Rev. **84** (1951) 1232
- [3] C. Itzykson and J.B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980)
- [4] H.-Y. Jin, C.-S. Huang and Y.-B. Dai, Z. Phys. **C56** (1992) 707; Y.-B. Dai, C.-S. Huang and H.-Y. Jin, Z. Phys. **C60** (1993) 527, Phys. Lett. **B331** (1994) 174; X.-H. Guo, H.-Y. Jin and X.-Q. Li, Phys. Rev. **D53** (1996) 1153
- [5] R. Akers et al., OPAL collaboration, Z. Phys. **C69** (1996) 195; Phys. Lett. **B353** (1995) 402
- [6] M. Anselmino, P. Kroll and B. Pire, Z. Phys. **C36** (1987) 89
- [7] G.P. Lepage and S.J. Brodsky, Phys. Rev. **D22** (1980) 2157; S.J. Brodsky, G.P. Lepage, T. Huang and P.B. MacKenzie, in *Particles and Fields 2*, edited by A.Z. Capri and A.N. Kamal (Plenum, New York, 1983), p. 83
- [8] X.-H. Guo and P. Kroll, Z. Phys. **C59** (1993) 567
- [9] E. Jenkins A. Manohar and M.B. Wise, Nucl. Phys. **B396** (1996) 38
- [10] M. Sadzikowski and K. Zalewski, Z. Phys. **C59** (1993) 677
- [11] M. Neubert and V. Rieckert, Nucl. Phys. **B382** (1992) 97
- [12] M. Neubert, Phys. Rev. **D45** (1992) 2451
- [13] J.L. Rosner, Phys. Rev. **D42** (1990) 3732
- [14] J.G. Körner and M. Krämer, Phys. Lett. **B275** (1992) 495

Figure Captions

Fig.1 The diquark-gluon-diquark vertex.

Fig.2 The weak transition diagram for $\Lambda_b \rightarrow \Lambda_c$.

Fig.3 The Isgur-Wise function for $\Lambda_b \rightarrow \Lambda_c(m_D = 700\text{MeV})$. For the solid line $\kappa = 0.02\text{GeV}^3$ and for the dotted line $\kappa = 0.1\text{GeV}^3$.

Fig.4 The differential decay width for $\Lambda_b \rightarrow \Lambda_c l \bar{\nu}$ ($m_D = 700\text{MeV}$). For the solid line $\kappa = 0.02\text{GeV}^3$ and for the dotted line $\kappa = 0.1\text{GeV}^3$.




Fig.4




Fig.1




Fig.2




Fig.3